

THE LOUISVILLE MEDICAL NEWS:

A WEEKLY JOURNAL OF MEDICINE AND SURGERY.

EDITED BY

LUNSFORD P. YANDELL, M.D., and L. S. McMURTRY, A.M., M.D.,

JOHN P. MORTON & CO., Publishers.

CONTENTS.

EDITORIAL—	PAGE.	ORIGINAL—	PAGE.
Malaria in Skin Diseases—A Correction,	253	A Case of Difficult Labor. By A. W. Reese, M.D.,	257
Post-Graduate Courses,	254	REVIEWS—	
MISCELLANY—		A System of Human Anatomy. By H. Allen, M.D.,	259
A Bequest,	255	MEDICAL SOCIETIES—	
A Dermatological Drama,	255	Pathological Society of Philadelphia,	261
Sad Outlook for the British Profession,	255	SELECTIONS—	
Gynecological Fortifications,	256	Treatment of Placenta Previa,	263
Case of Cobra Bite followed by Recovery,	256	Plica Polonica,	264
Guy's Hospital,	256	Operation for Hydrocele,	264
Brilliant Sentences from Britain,	256	Treatment of Soft Chancres and of Buboes by Salicylic Acid,	264
Sir Thomas Watson,	256	Syphilitic Polyuria,	264
Drs. Woodward and Barnes,	256		

NEW BOOKS

- ON SLIGHT AILMENTS; THEIR NATURE AND TREATMENT.** By Lionel S. Beale, M.B., F.R.S. Second, new, revised and enlarged edition.....Cloth, **\$1.95**
 "This new, revised edition of Dr. Beale's 'SLIGHT AILMENTS' is published simultaneously with the London edition by special arrangement between Dr. Beale and the American publisher. It therefore contains all the additions, illustrations, and a complete index, as in the London edition."
- THE NERVOUS SYSTEM.** A new and important addition to Appleton's popular series of "HEALTH PRIMERS." Square 16mo.....Cloth, **40 Cts.**
- A SYSTEM OF SURGERY; PATHOLOGICAL, DIAGNOSTIC, THERAPEUTIC AND OPERATIVE.** By Samuel D. Gross, M.D., LL.D. New, sixth edition, thoroughly revised, and greatly improved. Two vols.....Sheep, **\$15.00**
- PRACTICAL LABORATORY COURSE IN MEDICAL CHEMISTRY.** By John C. Draper, M.D., LL.D., Professor of Chemistry in the Medical Department, University of New York, and of Physiology and Natural History in the College of the City of New York. Sent by mail, postpaid, on receipt of price.....**\$1.00**
- A TREATISE ON THE SCIENCE AND PRACTICE OF MEDICINE, OR PATHOLOGY AND THERAPEUTICS OF INTERNAL DISEASES.** By Alonzo B. Palmer, M.D., LL.D. Part II.....Cloth, **\$5.00**
- A TREATISE ON THE PHYSIOLOGICAL AND THERAPEUTIC ACTION OF THE SULPHATE OF QUININE.** By Otis Frederick Manson, M.D., Professor of Physiology and Pathology in the Medical College of Virginia.....Cloth, **\$1.00**

JOHN P. MORTON & CO. have a full Stock of Medical Books always on hand.

AMERICAN PRACTITIONER.
Monthly, \$3.00 a Year.

JOHN P. MORTON & CO.
PUBLISHERS.

LOUISVILLE MEDICAL NEWS.
Weekly, \$3.00 a Year.

440 to 446 West Main Street, LOUISVILLE, KY.

THE LOUISVILLE MEDICAL NEWS and the AMERICAN PRACTITIONER will be sent to one address for one year to advance-paying subscribers for five dollars. Address publishers.

ISSUED EVERY SATURDAY.

THE IMPROVED
Harris Electro-Medical, Galvanic, and Faradic Batteries,
 PATENTED AND MANUFACTURED BY
A. C. HARRIS, Louisville, Ky.



FOR PHYSICIANS AND FAMILIES

Electro-medical appliances of all kinds supplied at reasonable prices.

Medical batteries of every description repaired at the shortest notice.

Good reliable agents wanted in all parts of the United States and Canada.

For illustrated catalogues, testimonials, price-lists, agencies, or any information concerning the batteries, address

G. T. CRAVEN & CO.,

General Agents for the United States and Canada.

141 and 143 Race Street, Cincinnati, O.

And 536 Third Street, Louisville, Ky.

QUALITY FIRST.

ESTABLISHED 1817.

ARTHUR PETER & CO.

LOUISVILLE, KY.

Importers and Wholesale Druggists,

MANUFACTURING PHARMACISTS.

Dealers in Druggists' Sundries, Choice Pharmaceutical Preparations. Original makers of Aromatic Elixir Grindelia, Aromatic Anti-constipation Elixir, Aromatic Elixir Licorice, Instruments of all kinds, Atomizers and Douches.

DEPOT FOR PARKE, DAVIS & CO.'S PREPARATIONS.

THE NEWCOMB-BUCHANAN CO.

DISTILLERS

AND

Wholesale Whisky Merchants

PROPRIETORS OF THE CELEBRATED

ANDERSON AND NELSON DISTILLERIES

Producing the very highest grades of Sour-mash and Fire-copper

BOURBON AND RYE WHISKIES,

34 Fourth St., LOUISVILLE, KY

JOHN P. MORTON AND COMPANY.

ESTABLISHED 1825.

BOOKSELLERS AND STATIONERS

MEDICAL
Books,

LAW
Books,

SCHOOL
Books,

PRAYER
Books,

HYMN
Books,

GIFT
Books,

JUVENILE
Books,

TOY AND
Miscellaneous Books.

Plain and Fancy Foreign and Domestic

STATIONERY

PENS (Gold and Steel)

PENCILS,

INKS,

INKSTANDS,

MUCILAGE,

PAPETERIES,

WRITING PAPER,

WRITING DESKS,

ENVELOPES,

ALBUMS,

POCKET BOOKS, etc.

ESTABLISHED 1825.

PUBLISHERS, PRINTERS, BINDERS

Child's First Book,
 Mother's Primer,
 Common School Primer,
 Davis's Pictures and Words
 Butler's Spellers,
 Butler's Readers,
 Butler's and Murch's
 Grammars,
 Bonnell's Composition,
 Towne's Mathematics,
 Nelson's Book-keeping,
 Butler's Speaker,
 Kavanaugh's Dramas,
 Huntoon's Speaker No. 1,
 Bronson's Elocution,
 Whately's Rhetoric and
 Logic,
 Peckham's Chemistry,
 Barker's College Chemistry
 Sedler's Chemical Experimentation,
 Barbee's Geology,
 Goodrich's History,
 Rogers's School Records,
 Barbour's Kentucky Digest
 Newman's Preading and
 Practice,
 Bush's Ky. Law Reports.

BOOK AND JOB

PRINTING

OF

All Kinds and Styles.

**RAILROAD,
 STEAMBOAT,
 MERCANTILE,
 COURT, and
 MISCELLANEOUS.**

**COURT RECORDS,
 JOURNALS,
 LEDGERS,**

And all kinds of

BLANK BOOKS

Made to order in handsome style on short notice.

S. H. KENNEDY'S**Concentrated Aqueous Extract**

—OF—

PINUS CANADENSISWHITE.DARK.**MEDICAL PROPERTIES,****MUCOUS ASTRINGENT.***A Most Valuable Remedy in all Diseases of the Mucous Surfaces
requiring an Astringent.*

NOTE TO PHYSICIANS.—I will take pleasure in forwarding you free a sample bottle sufficient to test fully its merits. A trial only is sufficient to establish its medicinal value.

—DIRECTIONS FOR USE.—

N. B.—The doses prescribed are for adults, for children the dose must be diminished according to their age.

AS AN INTERNAL REMEDY IN DIARRHŒA, DYSENTERY, NIGHT-SWEATS, HEMORRHAGE, OR PROFUSE EXPECTORATION. Mix one part of the extract with five of warm water; let it cool and take a teaspoonsful of the solution every three hours or oftener as the case may require.

For LEUCORRŒA and other VAGINAL DISEASES, dissolve a tablespoonful or two in a pint of warm water and inject twice a day. In obstinate cases a stronger solution should be used.

For PROTRUDING OR ITCHING PILES mix one-fourth of glycerine and apply as often as convenient; FISSURE OF THE ANUS, SORES, ULCERS, BURNS, or SCALDS, the extract should be applied in its full strength.

For CATARRH, dissolve a teaspoonful or two of the extract and a teaspoonful of salt in a pint of warm water and inject into the nostrils with a nasal douche twice a day.

For SORE THROAT, dissolve a tablespoonful of the extract in half pint warm water, let it cool, and apply as a gargle repeatedly during the day.

For GONORRŒA or GLEET, mix one part of the extract with three of water, inject two or three times a day.

For SKIN DISEASES, the extract should be used in its full strength, locally, and in obstinate cases a warm bath, in which a pint of the extract has been dissolved, should be taken several times a week.

To avoid STAINING OF LINEN the White P. C. should be used.

USED AND RECOMMENDED BY**J. MARION SIMS, M.D., New York.****ROBT. A. GUNN, M.D., New York.****R. WALKER, M.D., New York.****WM. RUSSELL, M.D., New York.****J. C. NIDELET, M.D., St. Louis.****T. F. RUMBOLD, M.D., St. Louis.**

And many other prominent Physicians throughout the country.

PREPARED ONLY BY**J. C. RICHARDSON, Chemist, ST. LOUIS, MO.**

Pinus Canadensis is prepared for use of Physicians only, and can be had from all reputable Druggists, wholesale and retail, in the United States, or from

RICHARDSON & CO., Wholesale Druggists, 710 North Main St., ST. LOUIS, MO.

TO PHYSICIANS.

BROMIDIA IODIA

FORMULA.—Every *fluid dram* contains 15 grs. EACH of pure Brom. Potas. and *purified* Chloral, and $\frac{1}{4}$ gr. EACH of *gen. imp. ext.* Cannabis Ind. and Hyoscyam.

DOSE.—One half to one *fluid dram* in WATER or SYRUP every hour until sleep is produced.

BROMIDIA is the Hypnotic *par excellence*. It produces refreshing sleep, and is *exceedingly* valuable in sleeplessness, nervousness, neuralgia, headache, convulsions, colic, etc., and will relieve when *opiates fail*. Unlike preparations of opium it does not lock up the secretions. In the restlessness and delirium of fevers it is *absolutely invaluable*.

The following physicians, having tested BROMIDIA, recommend it to the profession:

- EUGENE GRISSOM, A.M., M.D., LL.D., Raleigh, N. C.**
Superintendent North Carolina Insane Asylum.
- J. K. BAUDUY, A.M., LL.D., St. Louis, Mo.**
Prof. Nervous and Mental Diseases, Mo. Medical College.
- L. CH. BOISLINIERE, M.D., LL.D., St. Louis, Mo.**
Prof. of Obstetrics and Diseases of Women, St. Louis Medical College.
- WM. B. HAZARD, M.D., St. Louis, Mo.**
Prof. of General Pathology and Mental and Nervous Diseases, St. Louis College of Physicians and Surgeons.
- J. S. JEWELL, A.M., M.D., Chicago, Ill.**
Ed. "Journal of Mental and Nervous Diseases," and Prof. Nervous and Mental Diseases, Chicago Medical College.
- H. M. LYMAN, A.M., M.D., Chicago, Ill.**
Prof. Physiology and Diseases of the Nervous System, Rush Medical College.
- D. R. BROWER, M.D., Chicago, Ill.**
Ed. "Chicago Medical Journal and Examiner," and Prof. Nervous and Mental Diseases, etc., Woman's Medical College.
- I. N. DANFORTH, M.D., Chicago, Ill.**
Prof. Pathology and Diseases of the Kidneys, Woman's Hospital Medical College; President, and Lecturer on Pathology, Spring Faculty, Rush Medical College.
- D. D. BRAMBLE, M.D., Cincinnati, O.**
Dean; Prof. of Principles and Practice Surgery and Clinical Surgery, Cincinnati College Medicine and Surgery.
- WM. CLENDENIN, M.D., Cincinnati, O.**
Prof. Descriptive and Surgical Anatomy, Miami Medical College.
- J. B. MARVIN, M.D., Louisville, Ky.**
Prof. Chemistry, etc., and Clinical Lecturer on Nervous Diseases, Hospital College of Medicine.
- W. B. FLETCHER, M.D., Indianapolis, Ind.**
Prof. Physiology, Hygiene, and Clinical Medicine, Medical College of Indiana.
- W. J. SCOTT, M.D., Cleveland, O.**
Prof. Principles and Practice of Medicine, Medical Department Wooster University.
- H. H. POWELL, M.D., Cleveland, O.**
Prof. Obstetrics and Diseases Children, Cleveland Medical College.

Complaints have been made to us by physicians that some dishonest druggists *substitute an inferior preparation* made by themselves when BROMIDIA is prescribed. Physicians are cautioned to look out for these substitutions, because the lives of their patients may be endangered and their own reputation injured as well as ours. We have employed *detectives*, and shall protect our rights to the fullest extent of the law.

FORMULA.—Iodia is a combination of active principles obtained from the green roots of STILLINGIA, HELONIAS, SAXIFRAGA, Menispermum, and Aromatic. Each *fluid dram* also contains *five grains* IOD. POTAS. and *three grains* PHOS. IRON.

DOSE.—One or two *fluid drams* (more or less, as indicated) *three times a day*, before meals.

IODIA is the *ideal* alternative. It has been **LARGELY PRESCRIBED** in syphilitic, scrofulous, cutaneous, and female diseases, and has an established reputation as being the best alternative ever introduced to the profession.

The following physicians having tested IODIA, recommend it to the profession:

- EUGENE GRISSOM, A.M., M.D., LL.D., Raleigh, N. C.**
Superintendent North Carolina Insane Asylum.
- W. H. BYFORD, A.M., M.D., Chicago, Ill.**
President and Prof. Obstetrics, Woman's Hospital Medical College; Prof. Gynecology, Rush Medical College.
- R. M. KING, A.M., M.D., St. Louis, Mo.**
Prof. Physiology and Clinical Medicine, St. Louis College of Physicians and Surgeons.
- A. S. BARNES, M.D., St. Louis, Mo.**
Prof. Obstetrics and Diseases of Women, St. Louis College Physicians and Surgeons.
- C. D. PALMER, M.D., Cincinnati, O.**
Prof. Medical and Surgical Diseases of Women, and Clinical Gynecology, Medical College of Ohio.
- J. A. LARRABEE, M.D., Louisville, Ky.**
Prof. of Materia Medica and Therapeutics, and Clinical Lecturer on Diseases of Children, Hospital College of Medicine.
- M. F. COOMES, M.D., Louisville, Ky.**
Prof. of Physiology and Ophthalmology in the Kentucky School of Medicine.
- D. OVERLY CRIST, M.D., Indianapolis, Ind.**
Prof. Materia Medica and Therapeutics, Central College Physicians and Surgeons.
- N. W. WEBBER, M.D., Detroit, Mich.**
Prof. Medical and Surgical Diseases of Women, and Clinical Gynecology, Detroit Medical College.
- J. A. McCORKLE, M.D., Brooklyn, N. Y.**
Prof. Materia Medica and Therapeutics, Long Island College Hospital.
- J. M. BIGELOW, M.D., Albany, N. Y.**
Prof. Materia Medica and Therapeutics, Albany Medical College.
- J. L. WHITE, M.D., Bloomington, Ill.**
Ex-President Illinois State Medical Society.

BATTLE & CO., Chemists.

116 OLIVE STREET, ST. LOUIS, MO.

NESTLE'S MILK FOOD.

ITS PREPARATION AND VALUE FOR INFANTS.

Preparation.—For an infant under three months, mix one tablespoonful of the Food with ten of hot or cold water. Hold over the gas, lamp, or stove, with constant stirring, until it has boiled two or three minutes. Cool to about blood heat, and give in feeding-bottle. For a child three to ten months old, mix in proportion of eight of water to one of Food. For a pap, in proportion of five of water to one of Food.

Value.—Containing only Milk, Wheaten Bred Crust, and Sugar, this Food supplies all the elements necessary for complete alimentation, in the most easily assimilable form; the *Milk* furnishing Casein, Albumen, Hydrates of Carbon, and Sugar of Milk, while the *Wheaten Bread Crust* supplies Nitrogen, and is especially rich in *Saline Matter*, particularly in potash salts, mainly in the form of phosphates, and Carbon is obtained from the *Cane Sugar*. It makes pure blood, firm flesh, hard muscle, and tough bone. It is a sure preventive of Summer Complaint, and by its use the bowels can be kept in just the state desired. It is retained on the stomach often when everything else is rejected. The simplicity of its preparation and the uniformity obtainable are two points, the value of which can not be overestimated.

Particular Attention.—We do not claim that this Food will agree with all children. We do not think that any artificial food will ever be made which will do this, as nature sometimes fails, a mother's milk not agreeing with her own child. We only claim, what has been proved by its use for fifteen years past, that it will agree with a *larger proportion* of children than any other artificial food.

A pamphlet, by Prof. H. Lebert, of Berlin, giving fuller particulars of the Food, sent to any address on application to

THOMAS LEEMING & CO., Sole Agents,

18 COLLEGE PLACE, NEW YORK CITY.

For a perfectly pure CONDENSED MILK, free from starch of any kind, try Nestle's.

ew—370

MULTUM IN PARVO.

NO PRACTICING PHYSICIAN CAN AFFORD TO BE WITHOUT IT!

The American Practitioner's Simplified Visiting List and Account Book.

(COPYRIGHTED.)

This improved Visiting List and Account Book, which is so arranged as to be conveniently carried in the coat-pocket contains a NEW and SIMPLIFIED SYSTEM of keeping the accounts of practicing physicians with their patients, and which is so simple, complete, and accurate as to entirely dispense with the use of ALL other books, being complete and perfect within itself, and avoiding the necessity of posting or transferring the accounts, thus saving a great deal of writing time, and labor.

The book is 5 x 7½ inches (a good pocket size), handsomely and substantially bound in real Russia or Morocco, with Tuck, printed on bond paper.

No. 1600—Arranged for 200 Patients, Price, each	\$3 50
No. 1602 " " 300 " " "	4 50
No. 1604 " " 400 " " "	5 50

Sent to any address, postage paid, on receipt of price.

Full descriptive circular sent on application.

TERRELL, DIETZ & CO., Publishers,
514 W. Main St., Louisville, Ky.

MORTON'S POCKET SERIES.

No. 1.

DIET FOR THE SICK.

By J. W. HOLLAND, A.M., M.D.

Professor of Materia Medica, Medical Chemistry, etc.
University of Louisville.

Paper, 25 Cents; Handsomely bound in Cloth, 40 Cents.

"This book should be recommended by physicians to nurses and heads of families who have to provide meals for the sick."—*Virg. Med. Monthly*.

"It is quite readable and its conciseness commends it."—*Mich. Med. News*.

No. 2.

APHORISMS IN FRACTURE.

By R. O. COWLING, A.M., M.D.

Late Professor Principles and Practice of Surgery in the
University of Louisville.

Paper, 25 Cents; Handsomely bound in Cloth, 40 Cents.

They cover almost the entire field of fracture and present the subject in a tangible and practical shape such as has not heretofore been attempted. The Aphorisms will be found of a special service to students and general practitioners.

TO PHYSICIANS. LISTERINE.

FORMULA.—Listerine is the *essential Antiseptic* constituent of Thyme, Eucalyptus, Baptisia, Gaultheria, and Mentha Arvensis in combination. Each *fluid dram* also contains *two grains of refined and purified Benzo-Boric Acid*.

DOSE.—One teaspoonful *three or more times a day* (as indicated). As a local application to ulcers, wounds, and abscesses, or as a gargle, mouth-wash, inhalant, or injection, it can be used *ad libitum*, diluted as desired.

LISTERINE is a *powerful, safe, and pleasant Antiseptic*. The beneficial results following its use in Phthisis, Diphtheria, Catarrh, Dysentery, Scarlatina, Erysipelas, Smallpox, Typhoid and Malarial Fevers, etc., proves it to be a *restorative Antiseptic* of the very highest order of merit. It is the most efficient agent to disinfect the hands after surgical or gynecological operations, and is the *best injection* in Leucorrhœa, Gonorrhœa, etc. used in the proportion of from two to sixteen parts water, and one part Listerine.

Full Clinical Notes from the following and many other well-known physicians sent upon request:

PHILIP S. WALES,
Surgeon-General, United States Navy.

CHRISTOPHER JOHNSON, M.D.
Emeritus Professor of Surgery, University of Maryland, etc. etc.

MONTROSE A. Pallen, M.D., LL.D.
Professor of Gynecology, University of the City of New York, and Surgeon to the Maternity Hospital, etc.

HENRY O. MARCY, M.D.
Boston.

W. W. DAWSON, M.D.
Professor of Surgery, Medical College of Ohio, etc.

EDWARD W. JENKS, M.D., LL.D.
Professor of Diseases of Women, and of Clinical Gynecology, Chicago Medical College.

H. P. C. WILSON, M.D.
Ex-President Medical and Chirurgical Faculty of Maryland, and Baltimore Academy of Medicine; Vice-President American Gynecological Society.

OSCAR J. COSKERY, M.D.
Professor of Surgery, College of Physicians and Surgeons, Baltimore.

E. R. PALMER, M.D.
Professor of Physiology and Physical Diagnosis, University of Louisville.

HARVEY L. BYRD, A.M., M.D.
PRESIDENT; Professor of Obstetrics and Diseases of Women and Children, Baltimore Medical College.

JOHN A. OCTERLONY, A.M., M.D.
Professor of the Principles and Practice of Medicine, Kentucky School of Medicine.

E. B. STEVENS, A.M., M.D.
President Cincinnati Obstetrical Society.

E. H. GREGORY, M.D.
Professor of Surgery, St. Louis Medical College.

T. F. PREWITT, M.D.
DEAN; Professor of Surgery, Missouri Medical College, Surgeon to St. John's Hospital, etc.

P. V. SCHENCK, M.D.
Surgeon in charge St. Louis Female Hospital.

W. L. BARRETT, M.D.
Lecturer on Diseases of Women, St. Louis Medical College.

GEORGE J. ENGELMANN, M.D.
Professor of Obstetrics in the Post-Graduate School of the Missouri Medical College.

WM. PORTER, A.M., M.D.
St. Louis.

NATHAN S. LINCOLN, M.D.
Emeritus Professor of Surgery, Medical Department Columbia University, Washington, D. C.

FESSENDEN N. OTIS, M.D.
Clinical Professor Venereal Diseases, College of Physicians and Surgeons, New York City.

CHARLES T. PARKES, M.D.
Professor of Anatomy, Rush Medical College, Chicago.

PERCY NORCOP, M.D., F.R.C.S.
Formerly Surgical Dresser to Professor Lister.

JOSEPH TABER JOHNSON, A.M., M.D.
Professor of Obstetrics and Dis. of Women and Infants Med. Department University of Georgetown, D. C.

E. FLETCHER INGALS, A.M., M.D.
Professor of Diseases of the Chest and Physical Diagnosis, Rush Medical College, Woman's Medical College, etc. Chicago, Ill.

A. F. ERICH, M.D.
Professor Diseases of Women, College of Physicians and Surgeons, Baltimore.

THOMAS F. WOOD, M.D.
President Medical Society of North Carolina, Wilmington, N. C.

JAMES M. HOLLOWAY, M.D.
Professor of Surgery, Hospital College of Medicine, and Kentucky School of Medicine, Louisville, Ky.

DUNCAN EVE, M.D.
Professor of Surgery, Medical Department University of Tennessee.

A. M. OWEN, M.D.
Professor of Surgery, Evansville Medical College.

JOHN P. BRYSON, M.D.
St. Louis.

F. J. LUTZ, A.M., M.D.
Surgeon to Alexian Brothers' Hospital; Physician to Misericordia Asylum for the Insane and Nervous.

E. S. LEMOINE, M.D.
One of the Physicians to St. Luke's Hospital, St. Louis.

G. A. MOSES, M.D.
Lecturer on Clinical Gynecology, St. Louis Medical College.

J. B. JOHNSON, M.D.
Professor of the Principles and Practice of Medicine, St. Louis Medical College.

LAMBERT & CO., Manufacturing Chemists,
307 LOCUST STREET, ST. LOUIS.

 Listerine is sold by all Druggists on Physicians' Prescriptions. 

THE LOUISVILLE MEDICAL NEWS.

"NEC TENUI PENNĀ."

Vol. XIV.

LOUISVILLE, NOVEMBER 25, 1882.

No. 22.

LUNSFORD P. YANDELL, M.D., . . . }
L. S. McMURTRY, A.M., M.D., . . . } Editors.

MALARIA IN SKIN DISEASES—A CORRECTION.

Some time since the following paragraph appeared in the Michigan Medical News, and has been widely copied in the medical journals of the country:

A century ago John Hunter divided all skin diseases into three classes, one of which is cured by mercury and the iodides, a second by sulphur, and a third class which the devil himself can't cure. Dr. L. P. Yandell, who quotes Hunter as above, is given credit for a much less complex classification than even this. He attributes all skin eruptions to malaria. Quinine is a specific for malaria; ergo, quinine is the remedy for all skin eruptions. Q. E. D.

I trust that my confrères of the press will do me the kindness and the justice to publish the correction now given, as the matter is not only one of personal interest to the writer, but is of scientific interest to the profession. The subjoined extracts are from a supplement to a report read to the American Dermatological Association, September, 1877. A copy of this report will be gladly sent to any one desiring it:

"From the criticisms which have been made on my views, I find that I have not succeeded in making myself perfectly understood. What I have contended for, and what I have reiterated, is simply this: Malaria is the chief source of acute skin disease. Scrofula is the chief source of chronic skin disease. The more inveterate cases of skin disease are often due to the coexistence of these two things. The specific exanthems, of course, are not included here, but I con-

tend that their progress and termination are often largely influenced by the presence of malaria or struma. I do not claim that malaria and struma are the sole causes of the dermatoses. Indeed, many of the dermatoses may exist independently of malaria or struma, and most frequently some exciting cause is necessary to develop the cutaneous eruption. Among the exciting causes are irritants, injuries, insufficient or improper ingesta, vicissitudes of temperature, alcohol, dentition, menstruation, parturition, lactation, etc. The proofs of the truth of my views are, in the first place, that the diseases of the skin are cured more certainly and more quickly by the antimalarial remedies on the one hand, and by the antistrumous on the other, than can be done by any other line of therapeutics; and in the second place, that careful and painstaking investigation will, in the majority of dermatoses, make apparent the existence of the malaria or the struma, as the case may be.

"In conclusion, I desire to impress upon the reader that my views are not confined to the skin diseases. What produces disease here will produce it in all other organs of the body. What is true of dermatology is equally true of gynecology and ophthalmology and otology, and it is just as true of the diseases of all the other regions of the body."

Subsequent observation has confirmed my belief in the correctness of these views.

LUNSFORD P. YANDELL.

THE Hospital College of Medicine in this city will hold its sessions in the spring.

POST-GRADUATE COURSES.

In some remarks on this subject the Medical News, of Philadelphia, says:

In New York, the post-graduate idea has taken two directions. One school combines lectures with clinical and practical demonstrations; the other deals exclusively with clinical work. In a recent editorial, we decidedly commended the efforts made by our New York *confrères* to give practical direction to the two ideas of post-graduate instruction. But we are of the opinion that much of this method of instruction should be introduced into the regular sub-graduate curriculum, and that without the knowledge that is thereby obtained, a student is not fitted for graduation. The tendency of medical teaching now is to adopt this view and to supplement the didactic lectures with practical and clinical demonstrations in which the students themselves take an active part.

Recognizing the great need of practical eye, ear, and hand training, as a means for fitting the medical man for the work of his profession, Harvard and the two leading medical schools of Philadelphia have regular exercises in laboratory and hospital to secure this to each person. In every department—in clinical medicine, operative surgery, chemistry, materia medica, and physiology—each candidate for the doctorate is taught practically in the best sense of the word. At the Jefferson Medical School and at Harvard there is also an obstetrical laboratory, in which the principal obstetrical and gynecological operations on the female cadaver are performed in the regular curriculum, in the same way as on the living subject. At the University of Pennsylvania this course is embraced in the spring session, and we are informed that plans are being matured by which each student before graduation shall receive instruction in practical obstetrics at the bedside. . . .

In the methods inaugurated for the post-graduate instruction, there should be a combination of the clinical with the practical work in laboratory. Hence we fear that a merely clinical school will not accomplish the same good purpose.

New York has certainly taken the initiative in the organization of schools devoted solely to the instruction of graduates. Philadelphia has for several years provided suitable courses—as we have seen—for the same purpose, in connection with the existing medical colleges, and at the numerous special hospitals. It would also certainly be doing injustice to the Western schools if we failed to recognize the work which they are doing in the same way. Rush, the Medical College of Ohio, the University of Louisville, have provided excellent systems of post-graduate instruction, to which the profession has had access for several years. *Palmar qui meruit ferat.*

The post-graduate course of the University of Louisville follows immediately upon

the annual commencement, and furnishes to practitioners and recent graduates that practical and clinical instruction which is so essential for higher professional qualifications. Practical manipulatory instruction and laboratory work by which the eye, ear, and hand are educated is a part of the regular curriculum of study in this institution. The student, before graduation, is made familiar by actual practice, under the supervision of an instructor, with the use of the microscope, laryngoscope, ophthalmoscope, chemical manipulation, and the application of bandages, surgical dressings and appliances, as well as the use of the vaginal speculum, the stethoscope, and other aids to diagnosis and precision. The post-graduate course includes a higher grade of study, with advanced practical and clinical instruction, in order to furnish the practitioner and recent graduate those facilities formerly sought by American physicians in Europe, and which are essential to advanced skill and self-instruction in practical professional life. A post-graduate course is rapidly becoming a necessity in our educational system, and it is gratifying to see that our Eastern *confrères* appreciate the earnest and honest efforts made by the leading schools of the Southwest to give first-class facilities to students and practitioners. The announcement of the post-graduate course of the University of Louisville will appear early in the new year, and many practitioners throughout the Southwest will doubtless avail themselves of the advantages offered.

PROF. THEOPHILUS PARVIN.—The Edinburgh Obstetrical Society has just elected this distinguished teacher and writer an honorary member.

THE ARMY MEDICAL MUSEUM AND LIBRARY.—There is every reason to fear that at the next session of Congress an effort will be made to merge these valuable collections into the general Congressional Library.

MISCELLANY.

A BEQUEST.—The University of Louisville has lately received from the executors of the late Dr. George John Emil Renner five hundred dollars, bequeathed by the deceased to the dispensary department of the University.

Dr. Renner came to America soon after the Franco-Prussian war, through which he served in the Prussian army. He was a citizen of Freiberg, of excellent family, of high scholarship, proud, brave, and gentle.

He was graduated by the medical department of the University, March, 1877. He passed an exceptionally brilliant examination, and seemed destined to achieve the highest professional distinction. On the breaking out of yellow fever in Memphis in 1878 he hastened thither, one of the first volunteer physicians, and while laboriously engaged in the philanthropic work of ministering to the plague-stricken people fell a victim to the disease.

He died, as he had lived, loving God and his fellow-man, without fear of the future on either side of the grave, with a heart full of courage and kindness. His death was sorrowfully noticed at the time in these pages.

Deeply to be commiserated indeed is he who is without belief in a conscious existence beyond the grave, where such men as young Renner may again live and labor, carrying out there the pure and exalted purposes of a noble life broken off on hither side.

Let those who believe, pity and pray for those who are bereft of religious faith, rather than condemn them as malignant malefactors, as is the common custom of pious people.

Dr. Renner's means were limited, and for this reason his love offering to his Alma Mater, made on his death-bed, is all the more valued by the trustees and teachers of the University.

A DERMATOLOGICAL DRAMA.—The *Moniteur des Sciences Médicales et Pharmaceutiques* publishes an amusing "dermatological drama," called "King Sulphur," which is said to be played at the Hôpital St. Louis. Sulphur is King of Cutis, and has just conquered Acarus. He lays his crown at the feet of Queen Friction, who has aided him in the campaign, and implores her to become his honored queen. But she insists first on making an assault on Favus, and totally destroying his arrogant rule. If af-

terward Sulphur should burn with the same ardor, she will consent. Then she leads forth her army, attended by Axungia, while Sulphur marches in her train. Meanwhile the old tried generals, Hydrargyrum, Iodide of potassium, and Terebinth consult in angry conferences. Hydrargyrum is excited when he thinks that he, who has for forty years combated with so much glory all the forces of the Syphilides, should now be set aside for this Sulphur. Iodide laughs at his fears, and mocks at the silly tactics of Sulphur in such a war. Then we are introduced to the palace of Queen Eczema, wife of Herpes, who confides to her faithful attendant Acne her fears as to the future; she imagines she is losing her bloom, and is oppressed with vague fears. The news of the advance of Sulphur with Friction and Axungia causes vast alarm. Great preparations are made to resist him, but his attack is irresistible, and at length Favus, Eczema, Herpes, and all their generals have to acknowledge themselves vanquished by this terrible parasite and spore destroyer.—*Journal of Cutaneous and Venereal Diseases.*

SAD OUTLOOK FOR THE BRITISH PROFESSION.—During the past ten years we have watched the medical practice in London very closely, and we can truly say that, as regards general practice, it has been going from bad to worse, and from worse to still lower depths up to the present time. At no time has the outlook been worse than it is just now. Every department of practice is over-crowded, and a large proportion of practitioners have no notion whatever of professional honour and dignity. They act as if they were engaged in a trade, and do not care how small a fee they accept. There is scarcely a district in the metropolis but swarms with so-called "dispensaries" and open surgeries, where advice and medicine can be obtained for a shilling, and in the poor neighborhoods qualified surgeons are content with a fee of sixpence, whilst some are even content with fourpence for each attendance, and we have known a qualified doctor advertise to give *advice and medicine for twopence!* And it is of no use to complain of those who so degrade the profession; their answer is that they must accept low fees or starve.—*The Students' Journal.*

LINNÆUS taught that all diseases were produced by animalcula, or had an insect origin.

GYNCOLOGICAL FORTIFICATIONS.—The Medical Times thus alludes to Dr. Boze-man's paper, read before the American Gynecological Society: "The paper entitled 'Genital Renovation by Kolpostenotomy and Kolpoecpetasis in Urinary and Fecal Fistules,' is by Nathan Boze-man, M. D. It presents us in its first pages with the spectacle of the birth of a new word, for, following the example of the ophthalmologist, the gynecologist is now striving to bar the entrance to his specialty with mighty names. The reader of this paper must encounter and overcome cystostelosis, kolpo-kleisis, kolpostenotomy, kolpoecpetasis, kolpostenosis; and when there are thrown in a few other big words from outside sources, as pyonephrosis, etc., it is still more bewildering. Hysterocystokleisis is a good word, if it stops growing now. Then we have hysterokleidic, episio-kleisis, anakainosis. 'Some may object,' the writer says, 'to the introduction of so many new words;' but the introduction is a small matter, the difficulty is in recognizing them the next time one meets them in literary circles."

CASE OF COBRA BITE FOLLOWED BY RECOVERY.—Surgeon-Major Gunning (East India service) reports in the British Medical Journal, November 4th, this rare event. The patient was not seen for many hours after the injury. Surgeon Gunning is confident his treatment in no way affected the result. The snake was desquamating, and therefore not in a vigorous state of venom. Besides, he struck the victim's boot with his fangs repeatedly before he bit his hand, and thus doubtless expended the contents of the poison sac. The patient was ill and suffered decidedly, but was at no time in apparent danger. The cobra is the deadliest of all reptiles, and usually his bite results as in the following case described by Mr. Gunning: "A poor woman was bitten by a cobra about one hundred yards from where I am now writing, and was immediately carried across to me, but died in ten minutes after, or about thirty minutes after the infliction of the bite."

GUY'S HOSPITAL.—Dr. Steel, of Guy's Hospital, at a coroner's inquiry held by Mr. Payne the other day, opportunely explained that the reason why patients had frequently to be refused admission to that institution was simply because there was no room for them. Guy's and St. Thomas's are the only hospitals for the reception of accident cases in the South Metropolitan District, and as

the population is constantly increasing, without a corresponding addition in the ward accommodation at these hospitals being provided, they are unable to meet the growing demands made upon them.

BRILLIANT SENTENCES FROM BRITAIN.—Excerpts from Dr. Tripe's address in the Medical Times and Gazette of November 4th. Who is to blame, the speaker, the printer, or the editor? "Indeed, alterations in the proportion of land covered with water and forests have changed the climates of many countries to such an extent as to have rendered lands once fertile and healthy almost uninhabitable, or to have made unhealthy or sterile some localities which were formerly heathful and fruitful. . . . I suffered from these symptoms during the ascent of a mountain, at about ten thousand feet, and above, but they ceased on reaching the top and keeping myself still, showing that exertion had little to do with them."

SIR THOMAS WATSON.—American physicians will learn with profound concern that this revered physician and lovable gentleman has suffered an attack of paralysis indicative of cerebral lesions, and is gradually sinking. When his old pupil and devoted friend, Dr. George Johnson, reached his bedside, Sir Thomas, with calm demeanor and clear intellect, remarked, "This is the beginning of the end." Though the end is at hand, a noble career and the Lectures on the Principles and Practice of Physic will remain to testify the purity of character, superior professional attainments, and elegant scholarship which long ago won an envied and world-wide distinction.

DRS. WOODWARD AND BARNES.—It is with much regret that the profession learns of the continued impaired health of these well-known members of the profession. Dr. Woodward's health was not improved by his recent trip to Europe, and the condition of the ex-Surgeon General is such as to excite the greatest apprehension.

"I HAVE spoken almost exclusively of what I have myself seen and investigated." Dr. Cornil makes this remark in the preface to his work on Syphilis. If all authors would pursue this commendable plan we should have much fewer books as well as much smaller and better ones. The average medical treatise is chiefly composed of compilations from other bookmakers.

Original.

A CASE OF DIFFICULT LABOR.

BY A. W. REESE, M. D.

On Saturday, October 16, 1871, I was hastily summoned, at ten o'clock P. M., to the bedside of Mrs. L., of this place. On my arrival I found present in the sick woman's room Dr. H., the attending physician, Dr. W. C., called in consultation first, and Dr. S. P. added to the council previous to my arrival at the house. I discovered it to be an unusual case of labor. Dr. H. informed me that he had been called to the patient the preceding night (15th); that she had been in labor since two o'clock that morning; that she had just passed through two severe and alarming convulsions, and finally, that the medical gentlemen present all differed in regard to the presentation. Dr. H. gave it as his opinion that it was a presentation of the breech; Dr. W. C. thought it was the abdomen, and Dr. S. P. was sure that it was the shoulder of the fetus that presented itself to the touch.

Such then, in brief, was the history of the case, together with the differing views of the respectable medical men with whom I had been called to consult. The patient was profoundly under the influence of chloroform when I entered the room, Dr. H., the attending physician, administering the drug.

On concluding a statement of the above brief outlines of the case, Dr. H. asked me to make an examination and give my views, which I at once proceeded to do. The touch revealed a strange, unusually-shaped mass blocking up the entire pelvic cavity. In the course of a somewhat extensive obstetric practice I had hitherto met nothing like it. And yet, in spite of its seemingly anomalous character, I could not resist the conviction that the vertex was the presenting part. In fact I felt sure that it was the head that came in contact under the touch.

The shape of this cumbersome mass was, I admit, altogether unlike any other fetal head I had ever met before, and I must confess that this fact was rather against than in favor of my diagnosis in this knotty case. But on the other hand I was sure that I could feel the short, fine, silky hair that usually covers the fetal scalp.

Dr. S. P. could by no means agree with me in this diagnosis. He attributed the sensations derived by me from the touch, to abrasions of the cuticle upon the presenting

part of the fetus, as there had been a good deal of manipulation before my arrival in the case. Neither my judgment nor experience could approve this view of the matter.

Whilst making my examination, I rapidly reviewed, in my own mind, the conflicting and diverse opinions of my colleagues, and endeavored to determine their respective merits in a diagnostic point of view. First, then, I carefully scrutinized the position of Dr. H., the supposition of a breech presentation. And, truly, there seemed considerable grounds for his opinion. Here was a large mass, divided longitudinally, by a deep sulcus or groove, into two distinct, rather oblong hemispheres, which indeed bore a remarkable resemblance to the nates. But there were two features of the case which led me, unhesitatingly, to reject Dr. H.'s conclusion. In the first place, the hemispheres of the nates (if I may be allowed the expression) are soft, elastic, yielding, and pliable under the touch. In the present case these protuberances (whatever they might prove to be) were directly the reverse, being hard, dense, compact, inelastic, and, in short, felt to me just like bone covered by integument. Secondly, in passing my index finger from one end of this deep sulcus to the other (which, by the way, was no easy matter) I could discover neither the genital organs nor the anus; one or both of which must have been attainable in a breech presentation, except in a case of monstrosity, which latter is unusual and rare.

For these reasons I could not accept the view of the case under consideration as being a presentation of the breech. I next reviewed the opinion, expressed by Dr. W. C., that the abdomen was the portion of the fetus accessible to the touch. I endeavored to ascertain what features of the case could lead the Doctor's mind to this singular conclusion, for I had never met such a presentation, and was, moreover, skeptical as to its existence. The records of the profession sustain me in this opinion. Ramsbotham says, that in one hundred and fifty cases of transverse positions of the fetus, where he has been called to operate, he has met but *one case* of presentation of the abdomen. Chaillu denies their existence altogether; though in the American edition of that author's work, Dr. Gunning Bedford, the editor, mentions a case of the kind which he saw in consultation. Cazeaux, Dubois, and Naegele recognize but *two* trunk presentations, one for the right and

one for the left side. Madame Lachappelle denies the existence of such a presentation. This celebrated midwife declares that, in as many as forty thousand cases occurring at La Maternite, she had not met a *single case* of presentation of the abdomen.

In the case under consideration I could certainly expect to find either the soft, fluctuating, yielding parietes of the abdomen, the ensiform cartilage of the sternum, the symphysis pubis, or the insertion of the umbilical cord at the navel, if the belly, according to Dr. W. C., were the presenting part. But none of these portions of the fetus could be felt. I therefore excluded the abdomen from my diagnostic list. In confirmation of Dr. S. P.'s theory, that it was the shoulder, I could feel neither the axilla, any portion of the arm, the clavicle, fetal ribs, neck, acromion process, nor any other evidence that would lead me to the conclusion that it was a presentation of the shoulder, right or left.

On retiring for consultation I gave my opinion, and the reasons influencing my mind in entertaining the views expressed. Each of my colleagues, however, seemed "fully persuaded in his own mind" of the correctness of his own diagnosis. Such being the case, not much concert of action could be expected.

Finally, after much talk, it was agreed on to review the case, each of us to make another examination, and see what results could be obtained. We returned to the parturient chamber, and each one instituted a further examination, with the exception of Dr. H., the attending physician, who declined, stating that he was fully satisfied that it was the breech. Dr. W. C., the pioneer physician in the county, made a prolonged and rather tedious examination. He was succeeded by Dr. S. P., and lastly by the writer. We again retired for further consultation, and I must say that it was not a little amusing to see how positive each had become as to the correctness of his former opinion. No new light was thrown upon the subject.

Meantime it was becoming painfully evident that the vital forces of the patient were beginning to flag; the countenance was pale, the surface cool, the pulse feeble and growing quick and small, and the mind becoming despondent. These were serious symptoms, and showed that little time was to be lost in instituting some means for the woman's speedy relief.

The expulsive contractions of the uterus were powerful and continuous, but not a par-

ticle of advance was made by the presenting part of the fetus. As the result of further consultation resort was had to the forceps. Dr. S. P. volunteered his services in that direction, but after repeated and persistent efforts failed to deliver. After some time spent in these fruitless labors, the Doctor finally gave it up as a bad job, and asked me to try my hand. I took the handles of the forceps and withdrew the instrument from the patient's body. In reply to the Doctor's expressive look of inquiry, I said, "I am loth to use the forceps when there is room for doubt as to what part of the fetus they are to be applied."

I then made the third examination, as did also the two other consulting physicians, but without coming any nearer to an agreement than at first. I then made the suggestion that, regardless of the presentation, an effort should be made to reach the feet, and by turning the fetus deliver at once. This proposition met with general favor, for we had now reached a stage in the proceedings when any thing looking toward relief was gladly accepted. I was requested to make the attempt. I did so, and after great difficulty succeeded in reaching the feet, but found it impossible to turn. The two remaining consulting physicians both made similar efforts but without success.

At this stage of the case a final consultation was held, in which it was determined to use the perforator at once upon the most accessible part of the fetus, regardless of what it might ultimately prove to be, and thus by materially reducing its bulk, effect the speedy delivery of the woman, whose condition was now beyond question one of extreme peril. The perforator was therefore immediately brought into requisition. This procedure was instantly followed by an immense gush of water, a gallon at least in quantity, making its escape in a literal torrent. The blunt hook being then inserted into the opening made by the perforator, the fetus was speedily brought through the vulva. An inspection revealed a very large child with an enormously enlarged head.

The incision made by the perforator was directly in the center of the median line between the os frontis and the occiput, through the sagittal suture, thus putting the question of the presentation beyond all dispute. The sulcus felt by us was caused by the terrific pressure brought to bear by the uterine contractions upon the parietal bones of the fetal head.

On measurement, which was effected by

stuffing the cranial cavity with raw cotton and the use of a tape line, the proportions of the fetal head were found to be enormous. The distance from the nasal bones to the occipital protuberance was twenty-two and one half inches; circumference, measuring just above the ears, twenty-nine inches; from apex of chin to anterior fontanelle, nineteen inches. Unfortunately we neglected to weigh the child, but it could not have fallen short of fourteen or fifteen pounds. With the exception of its head it was well-shaped and healthy in appearance.

I saw this patient again on Sunday morning, the 17th, in consultation. Dr. W. C. and I called together. The symptoms were regarded as unfavorable. There was a pulse of one hundred and ten, inclining to be small and wiry, hypogastric tenderness, and pain and slight tympanitis. She was rational but restless and despondent. Dr. H. was still in charge of the case. I expressed my opinion to Dr. W. C., on our departure from the house, that she could not survive. The Doctor coincided in this view of the case. The prognosis proved correct, for she grew rapidly worse, and in a few days perished from metropéritoneal inflammation.

I am led to report this melancholy case, not for any purpose of self-glorification, not because I claim any special infallibility in diagnosis, or that I desire to appear "wise above that which is written," but because I think it an instructive case that may prove of some benefit to the profession, especially its junior members, and that there is sometimes "in a multitude of counsel" considerable confusion.

Accuracy in diagnosis is not always possible, even to the most experienced and skillful men. Mistakes will sometimes occur even with the best informed members of the healing art. I am satisfied that cases do arise where the wisest heads are sorely puzzled. Skill in diagnosis is the result of patient, laborious, and careful observation. Abernathy once said that "genius in a medical man consists in a patient observation of facts."

A man is a physician in the highest sense of the name, just in proportion to his knowledge of pathology and his skill in diagnosis, for "upon these two hang all the law and the profits." (Excuse the pun.)

The more we are impressed with this fact the more certainly shall we approximate that perfection in our noble profession which is the goal of our common ambition.

WARRENSBURG, MO., Nov., 1882.

Reviews.

A System of Human Anatomy, INCLUDING ITS MEDICAL AND SURGICAL RELATIONS. By HARRISON ALLEN, M. D., Professor of Physiology in the University of Pennsylvania. Illustrated with three hundred and eighty figures on one hundred and eighty-nine plates, many colored; also two hundred and fifty woodcuts. The drawings by Herman Faber, from dissections by the author. The whole in six quarto sections upon fine, thick paper. Philadelphia: Henry C. Lea's Son & Co. 1882.

To maintain a thorough familiarity with the essential details of anatomy is the most constant difficulty with which the practitioner of medicine contends. Regular annual review of this branch of study with dissection and demonstrations is usually impracticable in the midst of active practice, and no portion of medical knowledge so quickly fades into inaccuracy in memory as anatomical details. Many practitioners, in conscientious and intelligent recognition of the importance of this knowledge in connection with medical diagnosis and surgical procedures, endeavor to keep their knowledge fresh and accurate by occasional study and consultation of plates and drawings. These, unless very expensive, are usually inaccurate and unsatisfactory. Sibson's Medical Anatomy is an excellent work, but is limited to regional anatomy, with a brief and unsatisfactory text. MacLise's Surgical Anatomy is expensive and adapted only to the wants of the operative surgeon. Quain's large illustrated treatise is unwieldy and poorly illustrated. The well-known text books on anatomy are altogether descriptive, and do not treat of anatomy from the standpoint of either practical medicine or surgery. These latter works emphasize no especial feature, but treat of every detail in the descriptive method of elementary study. Teachers and students of anatomy have long felt the need of a work, thorough in details but conspicuous in important practical features; a work combining the advantages of faithfully depicted and artistically executed illustrations, together with a simple and practical exposition of the principles of human anatomy; a work suited to the wants of the student of medicine and surgery, and also to the general practitioner in the midst of his labors. It has fallen to the lot of our American confrère, Dr. Harrison Allen, of Philadelphia, to produce a work which fully meets these indications.

Human anatomy may be regarded an exhausted science, and many are the works

offered the profession on this subject, yet to write or teach anatomy successfully requires capabilities of the highest order. To teach medical and surgical anatomy with success and satisfaction necessitates a thorough practical knowledge of medicine and surgery acquired at the bedside. Moreover, in the preparation, arrangement, and practical adaptation of facts already known, as much ability and ingenuity may be displayed as in the search for the unknown in medical science. Dr. Allen undertook a severe task when he conceived the determination to produce a work on human anatomy which would meet the wants of both the student of medicine in his dissections and the practitioner amid the emergencies of a general practice. We know that it has required years of labor and diligent, patient, painstaking study and dissection to accomplish the task, and we congratulate him upon its completion.

The distinguishing feature of this work is that, while a thorough treatise on human anatomy, it is neither prepared from the standpoint of the scientist without knowledge of or sympathy with clinical requirements, nor from the standpoint of the surgeon, who often disregards the wants of the student and physician. The purpose has been maintained throughout of adapting the work to the wants of the student, the surgeon, and the physician.

The day is at hand when the American student of medicine will seek instruction in anatomy from the teacher who treats his subject from the standpoint of a practical medical man instead of from that of the scientist. We would not decry the study of anatomical science in its widest range, so attractive to the student of natural science and the very basis of all knowledge of living things. Human anatomy can never be appreciated without the study of comparative anatomy. No studies can offer such charms to the student of nature as are found in the study of anatomy and physiology in their widest domains. These studies, however, are for the few and not for the many. The student of medicine of the present day can not enter these inviting fields of study, but must perforce limit his investigations to the essential and practical. Hence we say, that teacher of anatomy will best meet the demands of the hour, who, thoroughly acquainted with his subject, has also a practical knowledge of medicine and surgery, and presents to his class in attractive and methodical manner, with due emphasis and abundant illustration, a knowledge of anatomy as applied to

medical and surgical practice. Dr. Allen is alive to these requirements, and has demonstrated the fact in the preparation and arrangement of his treatise on human anatomy.

The work is composed of six sections. The first is devoted to histology, and is prepared by Dr. E. O. Shakespeare, of Philadelphia, whose studies in both normal and pathological histology are familiar to the profession. The second section includes a consideration of the anatomy of the bones and joints. Both sections are admirably illustrated with plates of wonderful fidelity and beauty. The text is simple, concise, accurate, and lucid. The localization of diseased action and the surgical relationship are made prominent in connection with the consideration of every organ and part. The publishers have received the following note relative to the work from a well-known practical surgeon, who for many years publicly taught anatomy:

Gentlemen: Allen's Anatomy needs no commendation from me. It speaks for itself. I know of no similar work in any language which, either in illustration or text, is as well adapted to the wants of the medical student and the practitioner as this book of Professor Allen.

Yours truly,

D. HAYES AGNEW.

PHILADELPHIA, October 10, 1882.

This is high commendation, and the treatise fully merits it. The entire work will consist of six sections, arranged in portfolio form. This arrangement is most favorable for ready reference and convenience. The paper and typography are up to the well-known standard maintained by Messrs. H. C. Lea's Son & Co. It is sold only by subscription. The price is \$3.50 per section, the sections being delivered without additional cost and to suit the purchaser's convenience. The first two sections are now ready. Messrs. G. T. Craven & Co., 536 Third street, Louisville, and 141 Race street, Cincinnati, are the general agents for the West and South. Communications addressed to either office will elicit any further information desired in relation to Allen's Anatomy. M'M.

THERE is a true and a false medicine—the true consists in knowing how much we know; the false in pretending that all the arcana of disease and Nature is open to us. The true is noble and honest; the false is ignoble and dishonest.—*Thomas M. Dolan, F. R. C. S., in Med. Press.*

Medical Societies.

PATHOLOGICAL SOCIETY OF PHILADELPHIA.

The Pathological Society of Philadelphia met Thursday evening, October 9, 1882, the president, Dr. James Tyson, in the chair.

MYXOMATOUS TUMOR OF THE POSTERIOR CERVICAL REGION. Presented by Dr. Nancrede for Dr. W. G. MacConnell.

The tumor was removed by Dr. J. H. Brinton at the Jefferson College Hospital Clinic, some ten days since. The patient was a little boy aged four years, whose parents had first noticed the growth about two years ago. Latterly it has grown with considerable rapidity. It was of firm consistence, lobulated, and movable beneath the skin, giving the impression that it was a fibrous tumor. After removal, in addition to the above-mentioned characteristics, it was found surrounded by a capsule, and on section looked somewhat suggestive of myxoma; still it was thought by some to be merely a fatty tumor containing more fibrous tissue than usual.

Microscopic examination by Dr. MacConnell. Upon examining a frozen section stained with iodine, meshes of capillaries are displayed, in the walls of which the endothelial cells composing the vessels can be distinctly seen. The aforesaid meshes contain the mucoid structure traversed by large, pale, fusiform cells, the processes of which anastomose with each other. In addition many leucocytes are seen, and interspersing the growth in every direction numerous yellow, elastic fibres are readily distinguished.

When presenting the specimen, Dr. Nancrede commented on the rarity of such growths.

Dr. S. W. Gross said he had himself presented several gelatinous polypi of the nose, a number of years ago, which were most characteristic examples of myxomatous tissue. He could also recall a specimen of subcutaneous myxoma of the forearm, as well as the hematoid myxoma of the breast referred to by Dr. Nancrede. He was disposed to consider it the rarest of all neoplasms of the breast; indeed, he had never personally met with one, and when preparing his work on tumors of the breast he had written to numerous surgeons throughout the country, who all replied that they had never met with one affecting the breast.

Dr. Formad remarked that he had exhibited a myxomatous fibroma of the labium some years since, and said that the peculiar milky appearance assumed by the fluid when such growths were thrown into alcohol was a good diagnostic point.

Dr. Shakespeare said that his personal experience as to the rarity coincided with that of Dr. Gross. This specimen is one of the rarest forms, as most of the fibrillæ consist of yellow, elastic tissue. The rarity of myxomatous tumors seems to him to have much bearing on the views of Cohnheim and others as to the etiology of tumors. The observers insist that all tumors spring from the remains of fetal tissues, not made use of in tissue construction, which remain dormant in their embryonic condition until subjected to some irritation, when they develop into the various neoplasms. Now, tissue practically identical with

that found in myxomata pervades the fetus. How, then, is it that portions of this do not remain to give rise to myxomata? On the contrary, myxomata are among the rarest of the neoplasms.

BRAIN, LUNG, HEART, LIVER, SPLEEN, KIDNEY, AND BLADDER LESIONS. Presented by Dr. J. T. Eskridge.

The specimens showing the above lesions were removed from the body of a man aged sixty-eight years. The patient had become deaf in the right ear thirty years before, while suffering from some brain disturbance. Attacks of jaundice, with gradually-increasing permanent discoloration of the skin, had extended over a period of ten years. Since the early part of the year 1877 he had complained of incontinence of urine, an oppressed feeling over the hepatic region, dropsy in the feet and face, and a gradual loss of flesh and strength. The two years preceding his death he had been unable to work, but was only confined to bed five days. During the latter period his symptoms were in the order in which they were developed—great prostration, scanty secretion of urine, blindness for twenty-four hours preceding repeated convulsions, loss of speech, and almost total inability to swallow, although consciousness was preserved till coma ushered in the death scene. His temperature (axillary) did not rise above 100.5°. The surface head-temperature nearly equaled that of the axilla. No paralysis of the muscles of the face or extremities preceded death. The liver during life did not appear to be enlarged or altered in its outline.

The post-mortem examination revealed in the brain engorgement of the veins with some effusion, slight pia-mater inflammation in the neighborhood of the fissure of Rolando, apparent degenerative changes in the left island of Reil and anterior portion of the left tempero-sphenoidal lobe; in the pleura and lungs, old and numerous pleuritic adhesions, lobular and vesicular emphysema of the lungs, congestion of both lower lobes, and a nodule (probably cancerous) of the left apex; in the heart, fatty degeneration, dilated right ventricle, and incompetent mitral valves from ossific change; in the liver, multiple cancer without an increase in size or a nodular condition of the organ; in the spleen, marked increase of fibrous tissue and atrophy of the gland to one half or less its normal size; in the kidneys and ureters, the last stages of pyo-nephrosis, the glandular tissue being nearly destroyed, the pelvis were as large as a good-sized orange, and the ureters dilated so as to admit a man's thumb; in the bladder, great hypertrophy of the mucous membrane and decrease of the capacity of the viscus.

PRIMARY CARCINOMA OF PANCREAS AND LIVER. Presented by Dr. E. T. Bruen.

The interesting features pertaining to this case are the age of the patient (twenty-four years) and the rapidity of the abnormal processes. These rendered the diagnosis of malignant disease doubtful, till the appearance of nodular tumors in the liver. The family history was free from hereditary disease. The commencement of the disease dated from September, 1881. Death occurred on the 15th of January, 1882. At first the symptoms related solely to the digestive tract, such as dull and heavy sensation after eating, with acid eructations and

occasional vomiting. Subsequently sharp, cramp-like pains in the abdomen were a prominent symptom. After the lapse of a week there commenced general itching, and two weeks later the skin became yellow. This yellowness and itching never disappeared during the history of the case. At the autopsy the gall-duct was found obstructed by the enlarged head of the pancreas, so that extreme dilatation of the gall-bladder had ensued. This was probably the cause of the jaundice and not the liver disease itself. The bowels were regular and the appetite good when first seen. The case then presented evidences of partial obstruction of the gall-duct with digestive disorder, but without the symptoms characteristic of malignant disease of the stomach or bowels. By the middle of December, 1881, the liver-dullness then extended from the fourth interspace to three inches below the ribs; in the nipple-line from the ensiform cartilage the line of dullness extended to within one inch of the umbilicus. The hepatic region was tender upon pressure, especially over the epigastrium. The patient complained about this time of dull pain over the liver with griping pain in the abdomen. The pulse was ninety-six per minute. He had lost four pounds since admission, and looked thin. About this time a small inequality was noticed on the surface of the liver three inches above and a little to the inner side of the umbilicus. The spleen was enlarged. By January 7th the bosselation of the liver became distinct, and the enlarged gall-bladder, rendered irregular by gall-stones, presented a slowly-increasing, elastic, tender tumor, situated to the right of the epigastrium and umbilicus. By January 12th the pulse became rapid, one hundred and thirty per minute. The patient rapidly failed, and death occurred on January 15th.

Autopsy. The pancreas was enlarged to double its size, the growth chiefly occupying the head and compressing the common bile-duct. Microscopic examination showed it to be scirrhous carcinoma. The liver was thickly studded with nodules of medullary carcinoma, explaining the ante-mortem bosselated feel of the organ. The gall-bladder was distended to twice its normal size and contained a number of gall-stones.

Remarks. The duodenal end of the organ, as is usual, was the seat of the disease. In a paper on thirty-nine cases of primary carcinoma of the pancreas, in St. Bartholomew's Hospital Report for 1881, jaundice is stated to be always present, while in twenty-four cases of secondary carcinoma this symptom was noted in but seven cases. This is presumably from the secondary growth occurring in some other portion of the organ than its head. Murchison says that the characteristic symptoms of carcinoma of the pancreas are pain in the pancreatic region, sensible tinna, and persistent jaundice. To these Dr. Bruen would add intestinal dyspepsia, which differs in some essential features from the dyspepsia of organic disease of the stomach.

Dr. Musser remarked that he could vouch for there being a distinct tumor of the pancreas, as he was present at the autopsy. The case had been under his observation in the dispensary one month prior to admission to the hospital. On account of the age he was puzzled as to an exact diagnosis, although confident that the cause of the jaundice was obstruction. He noted among other symp-

toms the intense itching of the skin, a point of importance, Sims says, in the diagnosis of obstructive jaundice from that due to suppression. In five cases of tumor of the pancreas he had lately seen, all were accompanied by jaundice.

Dr. Bruen called attention to the uncertainty of bosselation as a symptom of malignant disease of the liver. He had presented to this society, only two weeks since, a liver exemplifying this condition in a marked degree, where nothing beyond cirrhosis in the stage of enlargement existed. The occurrence of carcinoma of the liver at so early an age is unusual, although Dr. Pepper had shown a specimen of the disease to this society some years ago, occurring in an infant.

Dr. Tyson said that there were two points in this case of great interest to him: First, jaundice in carcinoma of the pancreas, while it is a frequent symptom it is by no means an invariable symptom. Seven years ago he presented to this society the specimens from a case of primary pancreatic carcinoma where no jaundice had been present, and six months ago he presented to the society a specimen of enlargement of the head of the pancreas from a patient who also presented no symptoms of jaundice. Second, as to the diagnosis from cancer of the stomach he had noticed in his experience, as was mentioned in the history of Dr. Bruen's case, the absence of gastric symptoms. This negative symptom is of importance, since the tumor is often detected in precisely the same spot in both these diseases. The absence of gastric symptoms with intestinal indigestion, irrespective of fatty diarrhea, he considered the most reliable diagnostic points between carcinoma of the pancreas and stomach.

SPINDLE-CELLED SARCOMA OF THE SMALL INTESTINE. Presented by Dr. W. A. Edwards.

On September 23, 1882, I was asked to assist Dr. W. F. Atlee in the removal of an abdominal tumor. The patient, aged forty-eight years, whose menstruation had ceased at thirty-one years, first noticed the swelling in April last. On the day of operation she measured thirty-eight inches around the abdomen. The usual incision was made and the tumor reached, when its surface was seen to be of a dark purple hue, with a net-work of large veins ramifying in every part of its serous covering. A trocar and cannula were introduced, but nothing but blood followed the withdrawal of the trocar. The sac was then torn open, and its contents of a soft, brain-like consistence were emptied out. The growth was now turned out of the abdominal cavity. There was no distinct pedicle, but an attachment to the intestine of about the size of a half dollar was seen. Dr. Atlee says, "When I emptied the sac of its soft contents I examined carefully, with extreme care, the part fastened to the intestine, and my finger passed into the intestinal tube." A silk cord was tied around the attached portion and the remainder of the growth removed. The omentum was attached to the growth for a space of two inches. This was ligated and cut away, and the abdominal wound was closed, etc. Death occurred September 25th, at 4 A. M.

This growth sprang primarily from the submucous tissue of the small intestine and grew with great rapidity, as the patient was only aware of its presence last April, and by September she measured, as above stated, thirty-eight inches. Micro-

scopical examination of preparations taken from several portions of the growth clearly showed it to be a spindle-celled sarcoma, and a most typical one at that. The small intestine is an unusual site for this neoplasm. As far as I am able to ascertain there is no recorded instance of its occurrence in this situation. My friend, Dr. Formad, to whom I have shown the growth, concurs with me in this statement. On the day of operation I noted as well as I could the absence of all secondary deposits. The surrounding intestines and peritoneum were apparently normal, not even unduly hyperemic. No post mortem was permitted.

CHRONIC PARENCHYMATOUS NEPHRITIS COMPLICATING PHTHISIS PULMONALIS. Presented by Dr. James Tyson.

My object in showing these kidneys is to illustrate the morbid anatomy of the renal complications which so frequently attends the later stages of phthisis pulmonalis. It is very well understood that when edema of the feet and legs present themselves in cases of consumption the end is not far distant; but the renal complication which is at the bottom of such edema is often overlooked. It is of course not impossible that there should be edema in the last stages of phthisis, from simple alteration in the composition of the blood—a watery state of it—but in the majority of instances it means that the kidneys have become involved. As to the form of disease affecting the kidneys, it is acknowledged that it may be either lardaceous disease or chronic parenchymatous nephritis; but I think the impression prevails, at least it was my own until recently, that the amyloid kidney is the most frequent complication. I believe, however, that the chronic parenchymatous nephritis is more common, and it becomes a matter of interest, if not of importance, to be able to diagnose between these two conditions. It is well known that the microscopic and clinical characters of the urine in the various forms of kidney disease are often identical, so that no assistance is afforded by a study of the urine. The history of the case of course leads to neither particular form, but suggests both. One criterion only can I recall to aid us, and that is the presence of enlarged liver. So commonly associated is the enlarged amyloid liver with amyloid kidney that the absence of it almost necessarily indicates the presence of amyloid kidney. At least I am sure we would err less frequently if we were to consider all cases of renal diseases attending consumption unattended by enlarged liver to be parenchymatous nephritis rather than lardaceous disease. It is true we often have enlarged fatty liver in consumption, but the degree of enlargement never reaches that of the amyloid liver, and hereafter I shall be inclined to consider all cases of renal disease complicating consumption to be parenchymatous nephritis, unless they are associated with enlarged liver, when I shall conclude that they are instances of amyloid disease.

Dr. Bruen considered that the passage of large quantities of urine and a history of specific disease, or of prolonged suppuration preceding the kidney trouble, would warrant a diagnosis of amyloid renal disease.

Dr. Musser would ask whether the heart was hypertrophied, and what was Dr. Tyson's experience regarding hypertrophy of that organ in cases of amyloid disease, and of chronic tubal inflam-

mation of the kidneys. If not too late, he would like to call attention to the absence of cardiac hypertrophy, with an infinite degree of obstruction in the renal circulation, in the case Dr. Eskridge had presented. This is in direct opposition to the view held by some that the hypertrophy of the heart is a sequence of the renal obstruction in chronic interstitial nephritis.

Dr. Tyson replied that in this particular instance he did not see the heart, and could not tell whether it was hypertrophied or not. The same law holds good for amyloid kidney as for chronic nephritis; if the case last long enough, hypertrophy is sure to be found sooner or later.

ECCHYMOSES OF THE MUCOUS MEMBRANE OF THE STOMACH.—Presented by Dr. J. M. Barton:

The history of the case was that of chronic lung trouble. The stomach, upon being opened, presented an irregularly-shaped extravasation of blood about two thirds of an inch in diameter. The mucous membrane covering the effusion was healthy, as it was in the rest of its extent.

Dr. Tyson remarked that these effusions are not uncommon, but he had never seen them except in their pin-point form.

Dr. Roberts asked if there had been violent vomiting recently.

Dr. Barton replied that nothing of this sort had been observed for some months prior to death.

Selections.

Treatment of Placenta Previa.—(M. Hofmeier, Berlin): The author's conclusions and methods claim our attention on account of the excellence of his results. His material consisted of forty-six cases, thirty-five of which were delivered in one year, thus offering an excellent chance to judge of a method carried out by one man in so many cases. In judging of his method he first excludes from the forty-six cases three who were so far gone from hemorrhage when he arrived that there was no chance for any treatment. Of the remaining forty-three, in nineteen the location of the placenta was central, in sixteen lateral, and in eight marginal—a very large percentage of *central*.

The usual rule of treatment is to tampon until the cervix is sufficiently dilated. This rule the author opposes. He scarcely ever uses a tampon, and as to the cervix his rule is only to wait till clear symptoms of labor set in, either in uterine contractions or a funnel-shaped dilatation of the cervix. He then proceeds as actively and early as possible. This rule was followed in thirty-seven of the forty-three cases, after poor experience in other methods with the rest. In nineteen cases the cervix was partially dilated, in eighteen either entirely closed or with only a funnel-shaped dilatation. The earlier the operation the more of necessity is the choice limited to the combined external and vaginal version with one or two fingers, the Wigand-Braxton-Hicks method. This was done in thirty cases, the foot was brought down in three breech cases, three times internal version was performed and once the forceps applied. The combined turning was practiced as long as possible, and the hand introduced into the uterus only when absolutely necessary. The feet, having been guided to

the os, are seized, and by firm traction the buttocks effectually stop the hemorrhage. In cases of central position of the placenta, the author, in spite of all the arguments against it, is in favor of perforating the placenta and bringing the feet through. He did it in five cases, in three of which it was necessary on account of haste, and in two of which the child was already dead. It gives the mother the best chance, and the child's chance is by any method in such a case extremely small. The rest of the delivery, the author expressly states, should be *slowly* accomplished. The condition of the child may modify this rule, but even this must not make us increase the mother's risk. "One must have the courage to let a doubtful child's life be lost in his hands, rather than subject the mother to increased danger. The child is to be delivered *slowly*. Even so, the author's results were not bad as regards the children. Of thirty-seven, seventeen were already dead; of the twenty still living, six died (three premature and three from perforation of the placenta). Altogether, sixty-three per cent died and thirty-seven per cent lived, which is up to the usual standard. The statistics of the mothers, however, are much better. The author considers in them not only the immediate result, but the aftercourse of the case. In each case ergotin was given subcutaneously during extraction and the uterus was washed out afterward with a five per-cent carbol. sol.

Of the thirty-seven patients treated by these rules, one died. She had been treated for twenty-four hours by tampon, and the placenta was foul and offensive when the delivery took place, and she died seventeen days after from phlegmon and phlebitis of the thigh. H. believes she would have surely been saved if action had been prompter. This one case, out of thirty-seven, gives a mortality rate of 2.7 per cent, far ahead of any published rate, others having been 10 per cent, 16 per cent, and 40 per cent. After-hemorrhages occurred in some cases, but none which could not be controlled with ergotin, iced and hot-water injections. Of the six cases treated at an earlier date, and by the *waiting* method, one died; two had a long, severe lying-in; four children were dead. Of the whole forty-six cases, therefore, five died—10.8 per cent. The author adds two useful hints as to the location of the placenta. In nearly central location the smaller portion is on the side which is more loosened from the cervix lip. In placenta previa lateralis the proportion in favor of the right side is about 11.4.—*Ibid.*

Plica Polonica.—Dr. Ferdinand Lessing, of Winowa, Minn., relates the following case in the Medical Times of November 4th: Anna T., aged sixteen, went six weeks ago to the country with a lady friend, and, rambling about in the woods, they came to a cold spring and washed their feet in it. Next day A. felt chilly and languid, appetite impaired together with shooting pains through her limbs. A week after, she noticed that when combing her hair she could not pull the comb through as readily as heretofore, and by about a week more her hair was a matted mass. The symptoms of pains in her limbs had increased in proportion, as also a neuralgic pain in head and eyeballs. Two weeks before her death I was called, and found the sufferer in the following condition: She cried from the excruciating pain in limbs and head, the former being in a continuous state of tremor. Extremities cold, tongue clean, pulse sixty-five, appetite gone, insomnia complete,

and menstrual function stopped. I ordered her potassium bromide and chloral, also a tonic consisting of quinia sulph., iron, nux vomica, and arsenic. Gave her also wine and milk-punch *ad libitum*. The trembling of her limbs, as also the pain in head and eyes, had somewhat improved under this treatment in the course of a few days, yet her pulse grew weaker, and on the thirteenth day from the beginning she quietly passed away.

We once went a long distance, to Vienna, to see this disease, and were greatly disappointed to learn from Hebra that there was no such disease. He taught that so-called plica polonica was but an intense eczema occurring in broken-down constitutions, and was usually complicated and aggravated by lice. As to the hairs exuding a gelatinous matter, this is a physical impossibility. The exudation is from the scalp, as Hebra says, and merely clings to the hairs. This case of Dr. Lessings, however, is a rare and curious one, and well worthy of record.

Operation for Hydrocele.—Prof. W. T. Briggs, of Nashville, in a clinical lecture, published in the Nashville Journal of Medicine and Surgery, says on this point: Incision, as practiced originally by Sir Charles Bell, has of late fallen into disrepute, I being the only surgeon I am aware of who adheres to it as preferable to any other procedure. The operation consists of a free incision into the sac, its contents are allowed to flow off, and then place in the bottom of the wound between the walls of tunic a strip of lint. Acute inflammation soon follows. When it reaches the point of suppuration the lint is withdrawn, the walls of the cavity lying in contact adhere by adhesive inflammation and the cavity is thus obliterated. The wound heals by granulation, and the hydrocele is cured. This method is *always* effectual, and if conducted in the proper manner the inflammation is always under control. Prof. Briggs, of Nashville, availed himself of this case to urge the superiority of the operation by incision over any other. It is obvious, had this case been treated by injection, and an irritating fluid thrown into the sac, which was in such close relation with the testicle, disastrous consequences would certainly have followed. Recovery after the operation was a little tedious, but ultimately the patient was discharged entirely cured.

Treatment of Soft Chancres and of Buboos by Salicylic Acid.—1. The efficacy of salicylic acid in the treatment of soft chancres and of buboes appears to us to be unquestionable. While not an absolute specific, it is, in our opinion, capable of being most advantageously employed. 2. Odorless, only slightly painful in its application, soluble in alcohol and in glycerine, and leaving no stain on linen, it is preferable in these important respects to most other agents employed for the cure of the above-named affections, while perhaps inferior in certain other particulars to some among its rivals. 3. It may be resorted to in all cases, both when the sores are large and well-exposed, and when they are sloughing extensively, or are reached with difficulty; and it is equally available in private and in hospital practice. —*Four. of Cutaneous, Med. and Venereal Diseases.*

Syphilitic Polyuria.—Professor Semmola, of Naples, lately described this symptom in the *Revista de Ciencias Médicas* of Barcelona.

HARTER'S IRON TONIC.

FORMULA. Each dram of this preparation contains 1 grain of Iron, 2 grains Calasaya Bark, 1-200 grain Phosphorus, 1 grain Coca, 1 grain Viburnum, with a sufficient quantity of vegetable aromatics, Cologne Spirits, Sugar and Distilled Water.

HARTER'S IRON TONIC is a combination of Phosphorus, Calasaya Bark, Protoxide of Iron, Erythroxyton Coca, and Viburnum, associated with the vegetable aromatics in a pleasant and agreeable form, which has been so long a desideratum with the medical profession. It is pleasant and agreeable to the taste, having none of the inky flavors so peculiar to other preparations of Iron. In a low state of the system it will be found particularly efficacious. Iron restores color to the blood, and the Calasaya gives a natural healthful tone to the digestive organs. Phosphorus is a mild stimulant to the brain and nervous system, with especial action on the kidneys, bladder, and organs of generation, both in the male and female. The Erythroxyton Coca is a powerful nervous stimulant, through which property it retards waste of tissue, increases muscular strength and endurance, and removes fatigue and languor due to prolonged physical or mental effort.

TO THE MEDICAL PROFESSION.—We will take pleasure in forwarding you, free of charge, a sample bottle of the Iron Tonic, as a trial, which is sufficient to fully establish its medicinal value.

TESTIMONIALS.

F. FORCHHEIMER, M.D.

I consider HARTER'S IRON TONIC an excellent remedy, both in regard to its stomachic and general effects. The combination is very felicitous, and in my experience is always followed by good results.—Cincinnati, Aug. 4, 1882.

J. B. COX, M.D.

For Female Diseases HARTER'S IRON TONIC is par excellence. The Combination is well adapted to Anemia accompanied with Dysmenorrhea, the good results being attributed to the Iron, Phosphorus, and Viburnum.
St. Genevieve, Mo., Aug. 10, 1882.

J. S. DORSET, M.D.

I have been using HARTER'S IRON TONIC in my practice since 1875, and it has given me the most satisfactory results. I consider it a most excellent Tonic for general debility and nervous prostration.
Bonham, Texas, July 7, 1882.

DRS. RIPLEY & WALTON.

As a tonic for nervous prostration or exhaustion and want of vitality, HARTER'S IRON TONIC has never failed to do all that is claimed for it. It has given us complete satisfaction.
Philadelphia, Pa., July 5, 1882.

S. F. BARNES, M.D.

I have used Iron Tonic in a great variety of cases. From its admirable position its use is indicated in a wide range of diseases. It gives me excellent satisfaction.
Byron, Miss., Aug. 10, 1882.

J. S. FITZGERALD, M.D.

I am constantly prescribing Iron Tonic, it gives such general satisfaction. Where there is an opportunity it will reconstruct the most shattered and enfeebled constitution.
Butler City, Kansas, Aug. 15, 1882.

Recommended by many other physicians throughout the United States.

The Iron Tonic acts on the stomach and liver, increasing the appetite, assisting digestion, building up the weak, frail, and broken-down system, thereby making it applicable for dyspepsia in its various forms; loss of appetite, headache, insomnia, general debility, female diseases, want of vitality, nervous prostration or exhaustion, convalescence from fevers. It prevents impoverishment of the blood; is valuable in anemia, chlorosis, etc.

The Iron Tonic contains blood-making, force-generating, and life-sustaining properties, pre-eminently calculated to support the system under the exhausting and wasting process of disease, fevers, and other acute diseases, and to rebuild and recruit the tissues and forces, whether lost in the destructive march of such affections or induced by overwork, general debility in the most tedious forms of chronic diseases. It is friendly and helpful to the most delicate stomach. Does not cause nausea, constipation, or disarrange the digestive organs. Can be taken with impunity by the most delicate lady, infant, the aged or infirm, as by the sedentary student, whose system has suffered from over taxation of the brain; and where there is a fair remnant to build on, will reconstruct the most shattered and enfeebled constitution.

PREPARED BY

HARTER MEDICINE CO.

Practical and Analytical Chemists, ST. LOUIS, MO.

Harter's Iron Tonic is for sale by all Druggists on Physicians' Prescriptions.

FORTY-SIXTH ANNUAL ANNOUNCEMENT

OF THE

UNIVERSITY OF LOUISVILLE,

SESSION OF 1882 AND 1883.

FACULTY.

J. M. BODINE, M.D., DEAN.....	Professor of Anatomy and Diseases of the Eye and Ear.
LUNSFORD P. YANDELL, M.D.....	Professor of Principles and Practice of Medicine and Clinical Medicine.
E. R. PALMER, M.D.....	Professor of Physiology and Clinical Diseases of the Chest.
T. S. BELL, M.D.....	Professor of State Medicine and Sanitary Science.
JAMES W. HOLLAND, A.M., M.D.....	Professor of Pathology, Clin. Medicine, and Diseases of the Nervous System.
DAVID W. YANDELL, M.D.....	Professor of Surgery and Clinical Surgery.
THEOPHILUS PARVIN, M.D., LL.D.....	Professor of Obstetrics and Medical and Surgical Diseases of Women.
W. O. ROBERTS, M.D.....	Professor of Surgical Pathology and Operative Surgery.
JOHN A. OUTERLONY, A.M., M.D.....	Professor of Materia Medica, Therapeutics, and Clinical Medicine.
H. A. COTTELL, M.D.....	Lecturer on Medical Chemistry.
W. CHEATHAM, M.D.....	Clinical Lecturer on Diseases of Eye, Ear, and Throat.
L. S. McMURTRY, A.M., M.D., AND R. B. GILBERT, M.D.....	Demonstrators of Anatomy.

FEES.—Professors' Ticket, \$75.00; Matriculation Ticket, \$5.00; Practical Anatomy, \$10 00; Graduation, \$30.00 Hospital Ticket (required by the City), \$5.00.

SPECIAL AND OPTIONAL MANIPULATIVE COURSES.

H. A. COTTELL, M.D.....	Demonstrator of Microscopy.
B. BUCKLE, M.D.....	Demonstrator of Operative Midwifery.
W. CHEATHAM, M.D.....	Demonstrator of Ophthalmoscopy, Laryngoscopy, and Otoscopy.
L. S. McMURTRY, A.M., M.D.....	Demonstrator of Surgical Dressings.

The Spring Session of 1883 will open March 5th, and will continue until June 1st. It includes Clinical Teaching and Pharmaceutical work in the Dispensary, systematic recitations from Text-books, by a corps of examiners who have the use of the Museum for illustration, personal manipulations in Operative Surgery, Chemistry, Histology, Ophthalmoscopy, Laryngoscopy, and Otoscopy, under the supervision of Demonstrators.

The Spring Course is designed to be supplementary to the Regular Winter Course. Attendance upon it is voluntary, and does not count as a session.

The Fee for the Full Course is TWENTY-FIVE DOLLARS.

The Forty-Sixth regular Annual Session will commence on October 2, 1882, and will continue until March 1, 1883. Previous to this there will be a preliminary course of lectures free to all students, opening September 4th, and lasting until the beginning of the regular term.

The continued success of the practical exercises in Laboratories especially fitted with Beck's Microscopes, sets of Chemical Reagents, Manikins, Ophthalmoscopes, Laryngoscopes, etc., etc., has confirmed the wisdom of the Faculty in instituting these courses. Every facility and all needful apparatus will be furnished so as to make these teachings of permanent value to the student.

These special courses are optional. And it is recommended that first-course students should take Microscopy, for which a fee of \$5 will be charged, and second-course students the three other courses, for which a fee of \$10 will be charged.

It is urged upon all who seek to train their senses to the requisite degree of skill to make good diagnosticians and operators that at least one course of each of the manipulative branches be taken before applying for the degree.

CLINICAL MEDICINE AND SURGERY.

It is the determination alike of the Faculty and Trustees to secure to students that kind of information which will be most useful to them in active professional life, and it will be seen that no effort has been spared to make the University essentially a *practical and demonstrative* school.

The UNIVERSITY DISPENSARY, which is the property of the Faculty, affords great facilities to students. The building is upon the University grounds, and is open to patients and students throughout the year. It is the oldest institution of the kind in Louisville. It has obtained the confidence of the sick poor of the city, and its clinics are daily crowded with patients illustrating all varieties of disease. The advantages accruing to the University students from this source are among the chief attractions of the institution, giving them opportunities for attending cases and witnessing diseases in every phase. The Dispensary furnishes material for DAILY COLLEGE CLINICS from the following chairs: Clinical Medicine, Clinical Surgery, Diseases of Women and Children, Diseases of the Heart and Lungs, and Diseases of the Eye and Ear, Diseases of the Skin, and Diseases of the Nervous System.

In addition to the daily College Clinics mentioned, two Medical and two Surgical Clinics will be held weekly in the commodious amphitheater of the CITY HOSPITAL.

The Professors of Clinical Medicine and Clinical Surgery will lecture in the Hospital during the session. In addition to the above, the abundant clinical material of SS. MARY AND ELIZABETH HOSPITAL is at the command of the University Faculty.

FREQUENT EXAMINATIONS.

Universal experience has demonstrated the paramount importance of this mode of instruction as supplemental to lectures, and the Faculty has made a special provision for it. The wisdom of this action has been abundantly shown. The Faculty therefore devote additional hours for the purpose of a general "quiz," to be conducted by themselves.

Good boarding can be procured in the vicinity of the College at from \$3.00 to \$5.00 per week, fire and light included. Students on their arrival in the city by proceeding to the University, on corner of Eighth and Chestnut Streets, within three squares of the Louisville and Nashville Railroad Depot, will find the Janitor, who will conduct them to suitable boarding-houses.

A Post-graduate Course has been organized by the Faculty, which will follow immediately upon the winter session and continue six weeks. Special instruction will be offered to practitioners in various departments of medicine and surgery.

Address,

J. M. BODINE, M.D.,

Dean of the Faculty, Louisville, Ky.



BEEF PEPTONIDS,



A CONCENTRATED POWDERED EXTRACT OF BEEF, PARTIALLY DIGESTED AND
COMBINED WITH AN EQUAL PORTION OF GLUTEN.

We have pleasure in presenting, for the consideration of the Medical profession, "BEEF PEPTONIDS." We consider this product the most valuable that ever emanated from our Laboratory, and we feel confident it will be welcomed by the Profession in all parts of the world.

BEEF PEPTONIDS contains *only* the *nutritious* portions of the beef. It contains *no water* and *no inert matter* of any kind. We combine the dry Extract of Beef with an equal *portion* of Gluten to prevent a tendency to deliquescence, and in order to present the preparation in a powdered and portable form. It is well known that Gluten is the most nutritious substance found in the Vegetable Kingdom, and in nutritive elements is closely allied to Beef.

Four ounces of BEEF PEPTONIDS represents as much nutritive and stimulating properties as forty-eight ounces of the best lean Beef.

Four ounces of BEEF PEPTONIDS contains more nutritive elements than ten pounds of any extract made by Liebig's formula, and from four to six times more Albuminoids and Fibrinoids than any Beef Extract ever offered to the Medical Profession.

Our machinery and process for the production of BEEF PEPTONIDS are perfectly adapted to the *elimination* of all inert portions of the Beef, and the *retention* of all the nutritive constituents.

BEEF PEPTONIDS is *much less expensive* than any other preparation in the market, as it contains *neither water nor inert matter*.

The favor our preparation of BEEF PEPTONIDS received at the hands of DRs. AGNEW, HAMILTON, BLISS, KEYBURN, WOODWARD, BARNES, etc., the corps of eminent Physicians who employed the preparation with so much advantage in the treatment of the late PRESIDENT GARFIELD, proves conclusively its great value, not only as a food to be taken by the mouth, but also how important an agent it has been found in feeding by the Rectum.

Please refer to the very able article of Dr. D. W. BLISS in the New York Medical Record, July 15, 1882, in which he so frequently refers to BEEF PEPTONIDS having been used to so great an advantage, not only in the case of the late PRESIDENT GARFIELD, but many others as well.

We employ a reliable and experienced person to select the Beeves before they are slaughtered, and to superintend the killing and dressing. Great care is exercised in this respect, and none except the most healthy and suitable beeves are employed in making our BEEF PEPTONIDS.

Every physician will appreciate the importance of this care, for an Extract made from diseased Beef would not only be deleterious, but would, in many cases, produce lasting injury and fatal results.

The use of BEEF PEPTONIDS is indicated as follows:

Convalescence from all diseases, Fevers, Pneumonia, Weak Digestion, Diarrhea, Dysentery, Phthisis, Cholera Infantum, Marasmus, Sea Sickness, Excessive use of Alcoholic Stimulants; per Rectum in all cases where the stomach can not digest the food, and in debility resulting from any cause. Also a valuable adjunct in voyages and camp life.

We will be pleased to have the Profession every where test our assertions regarding this preparation, and for that purpose we will be happy to mail a sample to any regular practitioner desiring it; also circulars fully explanatory.

Thanking the Profession for generous support in the past, we
beg to remain,

Very respectfully,

REED & CARNRICK,

182 FULTON ST., NEW YORK.

ew-1y-408



Private Hospital for the Care and Treatment of the Insane.



CINCINNATI SANITARIUM.

The chief object of this institution is to furnish a retreat to that class of the insane whose education, social relations, and habits of life render indispensable more home-like comforts than can be commanded in the public asylums.

More than six hundred persons have already enjoyed the benefits of the institution, and fourteen States are at this time represented by one or more patients in our wards.

The location of the hospital is admirable in every respect. Situated in the midst of a highly cultivated country, five hundred feet above the Ohio River, the air is pure and wholesome at all seasons. The buildings are spacious, perfectly ventilated, thoroughly furnished, heated with steam, and lighted with gas.

A large library, pianos, billiard-table, a handsome hall for dancing, concerts, etc., afford ample means of recreation. In a word, all the means are provided calculated to arouse, elevate, and encourage the desponding or soothe, moderate, and tranquilize the excited.

Separate Department for nervous invalids and those suffering from the effects of opium and other narcotics. To Dr. O. EVERTS, formerly superintendent of the State Hospital for the Insane at Indianapolis, Ind., is confided the exclusive control of the medical and moral treatment of the inmates and the general management of the institution. The number of patients will be limited, so that each one will have the benefit of personal attention.

The cost of maintenance varies according to the rooms occupied and the special attention required. Special attendants are furnished to those who desire such extra service.

AL. P. COLLINS, *Secretary*.

60W-321

For further information, terms of admission, etc., address

Dr. O. EVERTS, Sup't, College Hill, Ohio.

TAKE ONLY AND INSIST ON "THE BEST OF AMERICAN MANUFACTURE."



PLANTEN'S CAPSULES*

Known as Reliable nearly Fifty Years.

*See note page 64, Prof. VAN BUREN AND KEYES on Urinary Organs.

PREMIUM FOR "GENERAL EXCELLENCE IN MANUFACTURE."



H. PLANTEN & SON, 224 WILLIAM ST., NEW YORK.

SOFT and HARD CAPSULES Filled, all Descriptions

EMPTY CAPSULES,

N. B.—All kinds of Capsules for mechanical purposes made to order, of any size.

New articles and private formulas a specialty.

Sold by ALL druggists.

SAMPLES FREE.



No. 00, Largest. No. 5 X, Smallest.

(Order by Number only.)

Boxes 100 each.

For taking medicines free of taste, smell, injury to the teeth, mouth, or throat. 100 (by mail), 50 cents.

RECTAL SUPPOSITORY (three sizes) CAPSULES. Box of 100 by mail, 50 cts.

HORSE CAPSULES (two sizes), ounce and half ounce. Box of 10 Capsules, 50 cts.

VAGINAL CAPSULES.

Capacity 10, 5, 3, 2, 1, ½, and ¼ Gr. Direct orders solicited.

SPECIFY PLANTEN'S CAPSULES ON ALL ORDERS. WE EMPLOY NO AGENTS OR TRAVELLERS.

THE CELEBRATED KENTUCKY ALUM SPRINGS WATER.

A Remedy for Diarrhea of Typhoid Fever, Summer Complaint, Diabetes Insipidus, Bronchitis, Whooping Cough, and very useful as a gargle in Sore Throat, a good wash for Ulcers, and excellent in Leucorrhœa and all other discharges from the mucous surfaces.

The following table shows the medical properties of this water as analyzed by Dr. L. D. Kastenbine, Professor of Chemistry in Louisville Medical College, and Professor of Chemistry and Practical Analysis in Louisville College of Pharmacy:

ANALYSIS.

Sulphate of Alumina.....	40.261
Persulphate of Iron.....	14.394
Sulphate of Iron.....	1.708
Sulphate of Lime.....	4.762
Sulphate of Magnesia.....	988
Chloride of Sodium.....	582
Silicic Acid.....	3.500
Sulphate of Potassa.....	Trace
Sulphate of Ammonia.....	Trace
Sulphate Manganese.....	Trace
Phosphoric Acid.....	Trace

Total per gallon..... 66.125

"I consider this a very pleasant Astringent and Tonic Mineral Water, especially adapted to chronic, gastric, and intestinal disorders, such as Dyspepsia, Diarrhea, and Dysentery.

April 12, 1882.

L. D. KASTENBINE, M. D."

R. A. ROBINSON & CO., Wholesale Agents,

Louisville, Ky.

Ask your Druggist for Pamphlet.

MALTINE.

MALTINE is a concentrated extract of malted Barley, Wheat and Oats. In its preparation the temperature does not exceed 150 deg. Fahr., thereby retaining all the nutritive and digestive agents unimpaired. Extracts of Malt are made from Barley alone, by the German process, which directs that the mash be heated to 212 deg. Fahr., thereby coagulating the Albuminoids and almost wholly destroying the starch digestive principle, Diastase.

LIST OF MALTINE PREPARATIONS.

MALTINE (Plain).
MALTINE with Hops.
MALTINE with Alteratives.
MALTINE with Beef and Iron.
MALTINE with Cod Liver Oil.
MALTINE with Cod Liver Oil and Pancreatine.
MALTINE with Hypophosphites.
MALTINE with Phosphorus Comp.
MALTINE with Peptones.

MALTINE with Pepsin and Pancreatine.
MALTINE with Phosphates.
MALTINE with Phosphates Iron and Quinia.
MALTINE with Phosphates Iron, Quinia & Strych.
MALTINE Ferrated.
MALTINE WINE.
MALTINE WINE with Pepsin and Pancreatine.
MALTO-YERBINE.
MALTO-VIBURNIN.

MEDICAL ENDORSEMENTS.

We append, *by permission*, a few names of the many prominent Members of the Medical Profession who are prescribing our Maltine Preparations:

J. K. BAUDUY, M. D., St. Louis, Mo., Physician to St. Vincent's Insane Asylum, and Prof. Nervous Diseases and Clinical Medicine, Missouri Medical College.

WM. PORTER, A. M., M. D., St. Louis, Mo.

E. S. DUNSTER, M. D., Ann Harbor, Mich., Prof. Obs. and Dis. Women and Children University and in Dartmouth College.

THOMAS H. ANDREWS, M. D., Philadelphia, Pa., Demonstrator of Anatomy, Jefferson Medical College.

B. F. HAMMEL, M. D., Philadelphia, Pa., Supt. Hospital of the University of Penn.

F. R. PALMER, M. D., Louisville, Ky., Prof. of Physiology and Personal Diagnosis, University of Louisville.

HUNTER McGUIRE, M. D., Richmond, Va., Prof. of Surgery, Med. Col. of Virginia.

F. A. MARDEN, M. D., Milwaukee, Wis., Supt. and Physician, Milwaukee County Hospital.

L. F. YANDELL, M. D., Louisville, Ky., Prof. of Clinical Medicine and Diseases of Children, University, Louisville.

JOHN. A. LARRABEE, M. D., Louisville, Ky., Prof. of Materia Medica and Therapeutics, and Clinical Lecturer on Diseases of Children in the Hospital College of Medicine.

R. OGDEN DORENUS, M. D., LL.D., New York, Prof. of Chemistry and Toxicology, Bellevue Hospital Medical College; Prof. of Chemistry and Physics, College of the City of New York.

WALTER S. HAINES, M. D., Chicago, Ill., Professor of Chemistry and Toxicology, Rush Medical College, Chicago.

E. F. INGALLS, A. M., M. D., Chicago, Ill., Clinical Professor of Diseases of Chest and Throat, Woman's Medical College.

A. A. MEUNIER, M. D., Montreal, Canada, Prof. Victoria University.

H. F. BIGGAR, M. D., Prof. of Surgical and Medical Diseases of Women, Homeopathic Hospital College, Cleveland, Ohio.

DR. DOBELL, London, England, Consulting Physician to Royal Hospital for Diseases of the Chest.

DR. T. F. GRIMSDALE, Liverpool, England, Consulting Physician, Ladies' Charity and Lying-in Hospital.

WM. ROBERTS, M.D., F.R.C.P., F.R.S., Manchester, England, Prof. of Clinical Medicine, Owens' College School of Medicine; Physician Manchester Royal Infirmary and Lunatic Hospital.

J. C. THOROWGOOD, M.D., F.R.C.P., London, England, Physician City of London Hospital for Chest Diseases; Physician West London Hospital.

W. C. PLAYFAIR, M.D., F.R.C.P., London, England, Prof. of Obstetric Medicine in King's College, and Physician for the Diseases of Women and Children to King's College Hospital.

W. H. WALSH, M. D., F.R.C.P., Brompton, England, Consulting Physician Consumption Hospital, Brompton, and to the University College Hospital.

A. WINN WILLIAMS, M.D., M.B.C.S., London, England, Physician Samaritan Free Hospital for Diseases of Women and Children.

A. C. MACRAE, M.D., Calcutta, Ind., Dep. Insp.-Gen. Hosp. Ind. Service, late Pres. Surg., Calcutta.

EDWARD SHOPPEE, M.D., L.R.C.P., M.R.C.S., London, England.

LENNOX BROWN, F.R.C.S., London, Eng., Senior Surgeon, Central Throat and Ear Hospital.

J. CARRICK MURRAY, M.D., Newcastle-on-Tyne, England, Physician to the N. C. H. for Diseases of Chest.

J. A. GRANT, M.D., F.R.C.S., Ottawa, Canada.

MALTINE is prescribed by the most eminent members of the Medical Profession in the United States, Great Britain, India, China and the English Colonies, and is largely used at the principal Hospitals in preference to any of the Extracts of Malt.

We will forward gratuitously a 1-lb. bottle of any of the above preparations to Physicians, who will pay the express charges. Send for our 28 page Pamphlet on Maltine for further particulars.

Address **REED & CARRICK,**

LABORATORY: Yonkers-on-the-Hudson.

189 Fulton St., New York.

Soluble Elastic Filled Capsules.

One of the most desirable improvements of modern pharmacy for the administration of medicine is the soft elastic capsule. These capsules, although so soft that the sides may be pressed together without injuring their walls, are so remarkable in their elasticity that they will immediately regain their original shape and size when the pressure is removed. This property renders them easy of deglutition, and they can be as readily swallowed as an oyster or the yolk of an egg. The composition of which they are manufactured is a compound of the finest French gelatine with glycerin, and is very soluble.

By employing the soft elastic capsule, the most nauseous dose is rendered slightly and palatable, for, covered by the capsule, its taste and appearance are completely disguised. Though the largest of them will contain a tablespoonful, capsules of this size are more easily swallowed than an ordinary pill. For this reason they are peculiarly adapted for the use of ladies and children. They are filled with castor oil, cod-liver oil, and other fixed oils, the various balsams, essential oils, ethereal extracts, etc., etc.

Our present list comprises sixty-seven formulæ, and is periodically revised, at which time additions are made of new compounds to keep pace with the demands of the profession.

The ingredients of capsules known as Nos. 6, 18, 21, 22, 24, 27, 34, 44, and 56, on our price list exist only in mechanical admixture. The heavy solid constituents, therefore, are merely held in suspension, and on standing they are deposited as an apparent sediment. This separation, however, does not impair the activity of the drug.

These capsules are made at an equable temperature, below that of extreme summer heat, and if subjected to a higher temperature and a moist atmosphere may adhere slightly to one another, or, especially in the case of the larger sized capsules, may become somewhat collapsed. These changes do not in the least impair the medicinal properties of the contents and do not imply any deterioration in the capsule itself. Such adhering or collapsed capsules regain their former condition in a cooler, lower temperature and drier atmosphere. These peculiarities are common to all soft or elastic capsules.

In ordering please specify our make of these capsules by writing the initials P., D. & Co. on your prescriptions. Only by doing so are you sure of obtaining the products of our laboratory.

Yours very truly,

PARKE, DAVIS & CO.

MANUFACTURING CHEMISTS,

DETROIT, MICH.

NEW YORK: 60 Maiden Lane and 21 Liberty Street.